



Captan 800WG

Fungicide for the control of certain diseases of fruit, greenhouse crops, ornamentals, turf and vegetables and some soil borne diseases as shown under Directions for Use.

<u>Active Ingredient:</u>	800g/kg captan
<u>Chemical Group:</u>	Cyclic Imide
<u>Formulation:</u>	Water dispersible granule
<u>Pack Size:</u>	10kg

Compatibility: Captan 800WG is compatible with most commonly used pesticides. It should not be combined with Bordeaux mixture, lime sulphur, oils or dicofof. Its use with emulsifiable concentrates is not recommended. If used in combination with sulphur on Kidds Orange, Delicious and some sulphur-shy varieties, some leaf spotting and drop may occur.

Captan 800WG is a proven fungicide, with broad-spectrum protectant activity against a range of diseases in many crops. Captan 800WG has a low risk of disease resistance, and is compatible with many IFP programmes.



Captan 800WG offers highly cost-effective disease control in a number of crops with many key benefits:

- Broad-spectrum **protectant activity** against a range of diseases ✓
- Low risk of resistance ✓
- A wide range of crop registrations ✓
- Highly cost-effective disease control ✓

NOTE: READ ENTIRE PRODUCT LABEL BEFORE USE. DIRECTIONS FOR USE:

CROP	DISEASE	RATE	CRITICAL COMMENTS
Apples, Pears	Black Spot, Glomerella, Sooty Blotch, Ripe Spot	75-125 g/ 100 litres water	Apply the high rate every 7 days from greentip through to petal fall. Captan products may increase the risk of Russet on some varieties. Where this is of concern use alternative materials between full bloom and second cover. Where Glomerella, Ripe Spot or Sooty Blotch are a problem use the high rate from January to harvest. Do not apply to Anjou Pears.
Blueberries	Stem Blight, Leaf Spot, Botrytis	125 g/ 100 litres water	Spray just before flowering; at 10% flowers open; at petal fall; and every 10-14 days until harvest.
Boysenberries, Loganberries, other Brambles	Canespot		Apply at petal fall and repeat every 10-14 days until harvest.
Citrus	Brown Rot	125-240 g/ 100 litres water 125 g/ 100 litres water	As a skirt spray before winter rains. Thoroughly wet trees to a height of 1 metre. Repeat 10 weeks later. OR Spray entire tree in autumn and repeat 4-6 weeks later. Do not use oil products within 14 days either side of Captan 800WG.
Red and Black Currants	Leaf Spot, Botrytis	125 g/ 100 litres water	Apply every 10-14 days from green tip until harvest.
Gooseberries	Leaf Spot, Botrytis	125 g/ 100 litres water	Apply just before flowering; again at petal fall; pre-harvest and post-harvest. For Botrytis Flower Rot, application at 5-7 days intervals over flowering may be necessary, depending on the weather.
Grapes	Dead arm, Anthracnose, Botrytis	125 g/ 100 litres water	Apply every 14 days, commencing when shoots are 5-8 cm long.
Raspberries	Cane Spot, Sour Blight, Botrytis	125 g/ 100 litres water	Main Crop: Apply every 10-14 days from greentip until harvest. Autumn Crop: Apply every 10-14 days from November until harvest. Thorough coverage is essential.
Stonefruit	Brown Rot, Scab, Shothole	125 g/ 100 litres water	Apply every 7 days from bud movement until bloom, then at 3-4 day intervals until petal fall, then fortnightly until 3 weeks before harvest. Apply weekly until harvest.
Strawberries	Leaf Spot, Botrytis	125 g/ 100 litres water or 3 kg/ha	Apply as necessary for Leaf Spot. For Botrytis apply at 7 day intervals from the beginning of flowering and between picking if wet conditions prevail.
Peas, Beans	Root Rot, Damping Off	7-9 kg/ha	Broadcast preplanting and work into upper 10 cm of soil.
Carrots	Ring Spot, Cercospora Blight	3 kg/ha	As necessary.
Cucurbits	Anthracnose	1.5-3 kg/ha	As necessary.
Lettuce	Leaf Spot	4.5 kg/ha	Every 10-14 days from rosette stage. The addition of a sticker is recommended.

SEED TREATMENT

CROP	RATE	CRITICAL COMMENTS
Peas	100 g/ 500 mL water/ 100 kg seed	Keep stock solution well agitated and do not allow to stand overnight. Captan 800WG does not contain a sticker. A suitable sticker should be added to the stock solution. Contact your local representative for further advice. Ensure treated seed is clearly labelled.
Potato seed	125g/100 litres water	For control of Seed rot and Damping off use as a pre-planting dip within 6 hours of cutting

WITHHOLDING PERIODS:

Blueberries, Currants, Gooseberries, Raspberries, Strawberries: DO NOT APPLY LATER THAN 1 DAY BEFORE HARVEST

Cucurbits: DO NOT APPLY LATER THAN 3 DAYS BEFORE HARVEST.

Lettuce, Stonefruit: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

Boysenberries, Loganberries, other Brambles, Carrots, Pipfruit: DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.

Grapes: DO NOT APPLY LATER THAN 21 DAYS BEFORE HARVEST.

Citrus: DO NOT APPLY LATER THAN 28 DAYS BEFORE HARVEST

Export crops: Consult your exporter for advice about any restrictions

Registered to and Distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 1250, NEW ZEALAND.

Ph: +64-9-412-9817; Fax: +64-9-412-9807

www.adriacp.co.nz

Captan 800WG is Registered pursuant to the ACVM Act 1997, Approval No. P7751