



Curator

For the control of certain diseases in mushrooms, ornamentals and stonefruit.

<u>Active Ingredient:</u>	462g/litre prochloraz as the manganese complex.
<u>Chemical Group:</u>	Imidazole
<u>Formulation:</u>	Wettable Powder
<u>Pack Size:</u>	1 kg
<u>Compatibility:</u>	Curator is compatible with most commonly used fungicides and insecticides. DO NOT use with alkaline materials such as lime sulphur or Bordeaux mixture.

Curator is a new high-quality formulation of '*prochloraz*', and is the first generic competitor to Octave®¹ to be registered in NZ.

Curator is a proven performer in mushrooms, ornamentals and stonefruit, and controls a wide spectrum of diseases. Curator is non-systemic but has strong contact activity and translaminar mobility.



Curator offers highly cost-effective disease control, with many key benefits:

- Excellent **activity** against *Verticillium spp.* In **mushrooms** ✓
- **Broad spectrum** disease control in **ornamentals** with a high degree of **crop safety** ✓
- **Good tank-mixing flexibility** ✓
- **Reliable performance** and highly **cost-effective** disease control ✓

GENERAL INFORMATION:

Curator is a DMI fungicide which may be applied as a spray treatment to ornamentals and stonefruit, as a dip treatment to ornamentals, and a drench treatment in mushrooms.

RESISTANCE MANAGEMENT: Resistance to this and other DMI products could develop from repeated use. To minimise this risk use strictly in accordance with label instructions.

DIRECTIONS FOR USE:

The method of application shall be limited to ground based application only.

Mixing: Thoroughly pre-mix the required amount with a little water to form a slurry. Add water whilst agitating. For spray treatments maintain agitation while mixing and applying. Also agitate thoroughly after a stoppage, before recommencing spraying. Use immediately after mixing.

COMPATIBILITY: Curator is compatible with most commonly used insecticides and fungicides. DO NOT use with alkaline materials such as lime sulphur or Bordeaux mixture.

RECOMMENDATIONS: Rates suggested are for high volume spraying to run-off. For concentrate spraying, adjust dilution rate accordingly.

CROP	DISEASE	RATE	REMARKS
Mushrooms	Brown spot (<i>Verticillium spp.</i>)	3g/square metre in 1 litre of water as a drench treatment.	Apply 7-9 days after casing. Do not apply within 10 days of picking. Apply a maximum of 3 applications per season.
Ornamentals	<i>Colletotrichum spp.</i> , Glomerella leaf spot, <i>Monochaetia spp.</i> , <i>Pestalotia spp.</i> , Powdery mildew	100g/100 litres in sufficient water to obtain thorough spray coverage. 100g/100 litres as a dip.	Apply at first signs of infection and repeat at 14 day intervals. Immerse whole plants in dip treatment prior to shipping. Apply a maximum of 3 applications per season.
Stonefruit	Brown rot	25g/100 litres	Apply at pink and repeat at 7-10 day intervals until petal fall. Shorter intervals may be required in wet weather. Use in mixture with non-DMI fungicides and apply no more than three treatments per season. Apply a maximum of 3 applications per season.

WITHHOLDING PERIODS:

Mushrooms: 10 days

Stonefruit: Do not use after petal fall

Registered to and Distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 1250, NEW ZEALAND.

Ph: +64-9-412-9817; Fax: +64-9-412-9807

www.adriacp.co.nz

Curator is Registered pursuant to the ACVM Act 1997, Approval No. P7738

®1 Octave is a Registered Trademark of Bayer AG and is registered pursuant to the ACVM Act 1997, Approval No. P3055