



# Ultima<sup>®</sup> 240EC

For the selective control of Wild oat, Lesser Canary grass and Gnawed Canary grass in Wheat, Triticale, Ryecorn and Durum wheat.

<u>Active Ingredient:</u>	240g/kg clodinafop-propargyl
<u>Chemical Group:</u>	Carboxylic acid derivative
<u>Formulation:</u>	Emulsifiable concentrate
<u>Pack Size:</u>	1 Litre
<u>Compatibility:</u>	Ultima is compatible with emulsifiable mineral oils approved for use on cereals. The addition of other herbicides is NOT recommended. Broadleaf herbicides should be applied 10 days before or after Ultima.

Ultima<sup>®</sup> is a selective post-emergent herbicide which provides excellent, cost-effective control of wild oats, lesser canary grass and gnawed canary grass in wheat, triticale, ryecorn and durum wheat.



Ultima<sup>®</sup> offers highly cost-effective wild oat and phalaris control:

Excellent crop safety ✓

Rainfast in 2 hours ✓

Flexible application timing ✓

Prevents further weed growth within 48 hours of application ✓

## GENERAL INFORMATION:

Ultima is rapidly absorbed by the leaves and shoots of grass weeds and is translocated to the growing point preventing further growth within 48 hours after application. DO NOT apply Ultima if rain is expected within 2 hours after application.

## DIRECTIONS FOR USE:

**Volume of application: Apply Ultima in a minimum of 200 litres of water/hectare.**

Crop	Weeds controlled	Weed stage	Rate/ha	Critical comments
Wheat, Ryecorn, Triticale, Durum wheat	Wild oat	3 leaf to 1 <sup>st</sup> node	100ml plus 1 litre mineral oil	
	Lesser Canary grass (Phalaris minor)	3 leaf to 4 tillers	100ml plus 1 litre mineral oil	
	Gnawed Canary grass (Phalaris paradoxa)	3 leaf to 3 tillers	100ml plus 1 litre mineral oil	Long term control is enhanced by good ground cover from an actively growing crop.

## EQUIPMENT:

Ultima can be applied using conventional boom spraying equipment and is non-corrosive.

## COMPATIBILITY:

Ultima is compatible with emulsifiable mineral oils approved for use on cereals. The addition of other herbicides is NOT recommended. Broadleaf herbicides should be applied 10 days before or after Ultima.