



Valeo[®] 500SC

A FUNGICIDE for controlling diseases of fruit and roses.

<u>Active Ingredient:</u>	500g/litre folpet
<u>Chemical Group:</u>	Cyclic Imide
<u>Formulation:</u>	Suspension concentrate
<u>Pack Size:</u>	5 Litre
<u>Compatibility:</u>	Valeo 500SC is compatible with most commonly used insecticides and fungicides. Do not mix with Bordeaux mixture, lime sulphur or oils.

Valeo 500SC is a protectant fungicide, with activity against a range of diseases. Valeo has a low risk of resistance, and has no cross-resistance to most commonly used fungicide chemical groups.



Valeo[®] offers highly cost-effective disease protection - with many key benefits:

Strong protection against Verucosis and melanose in citrus ✓

Proven protection against Black spot in roses ✓

Good alternating chemistry in Botrytis control programmes ✓

Broad spectrum ✓

MIXING AND COMPATIBILITY: Pour the measured quantity of Valeo 500SC into the partly filled spray tank, then top up with water. Agitate thoroughly during filling and spraying.

Valeo 500SC is compatible with most commonly used insecticides and fungicides. Do not mix with Bordeaux mixture, lime sulphur or oils.

It is an offence to use this product on animals.

DIRECTIONS FOR USE:

APPLES: Black spot, glomerella – use 150-200ml per 100 litres. Apply at 10-14 day intervals, commencing between mid-December and early January.

BUSHFRUIT: (blueberries, currants and gooseberries) and canefruit (blackberries, loganberries, raspberries and youngberries): Botrytis, leaf spots – apply 150-200ml per 100 litres of water over flowering. Repeat at 7-14 day intervals from flowering until harvest. Use the higher rate when conditions favour infection.

CITRUS: Verrucosis, melanose – apply 250ml per 100 litres prior to blossoming and repeat when flowering has finished.

GRAPES: Anthracnose, downy mildew – use 200ml per 100 litres of water at regular intervals from budburst.

Dead arm- apply 200ml per 100 litres of water after budburst when shoots are 3cm long.

Note: Application closer than 3 weeks before harvest may delay fermentation in wine making.

ROSES: Black spot, powdery mildew – apply 200ml per 100 litres of water at regular intervals from budburst.

IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

WITHHOLDING PERIOD: All crops – 3 days.

RESISTANCE MANAGEMENT: Valeo 500SC is a cyclic imide fungicide. Resistance to Valeo 500SC and related fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label instructions. Apply a maximum of 4 applications in one growing season.

Registered to and Distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.
Ph: +64-9-412-9817; Fax: +64-9-412-9807

www.adriacp.co.nz

Valeo 500SC is registered pursuant to the ACVM Act 1997, Approval No. P8158